

NEUROLOGICAL VISION SCREENING AFTER TBI CLINICAL PRACTICE GUIDELINE

The following are recommendations for a thorough screening of neurological vision impairment, based on the current scientific literature, which can then be used to develop a client-centered treatment plan, in collaboration with an OD or MD (vision specialist). A visual screener can be completed by a trained, non-vision specialist clinician, but it does not substitute for a full evaluation by a vision specialist. It allows us to identify appropriate candidates, make needed referrals, and opens up the line of communication between the therapist and vision specialist. It is not within an OT's scope of practice to diagnose. Many conditions can be treated by a licensed therapist who has been trained in vision therapy techniques, but must be in collaboration with a vision specialist. This collaboration allows integration of assessments of the eye/lower-order cerebral mechanisms (visual function) and person-level dysfunction/higher-order cerebral mechanisms (functional vision), targeting quality of life.³⁷ The ideal setting for providing the patient with brain injury with appropriate vision care and therapy is a team of providers working together in a vision clinic.³⁶

Vision Specialists

- Optometrist – Doctors of Optometry (OD) diagnose and treat vision problems, eye diseases and related conditions, prescribe eyeglasses and contact lenses and provide medication to treat eye disorders. They cannot perform surgery but often provide pre and post-surgical care.
 - ***It is important to refer neuro/concussion patients to an Optometrist who can provide a neuro-optometric/sensorimotor evaluation, as not all Optometrists perform this type of assessment. (See specialists on page 10).**
- Ophthalmologist – MD who specializes in the medical and surgical care of the eyes and visual system and in the prevention of eye disease and illness.
- Neuro Ophthalmologist – Subspecialists of neurology and ophthalmology who address visual problems related to the nervous system.

www.aoa.org

<http://www.eyesitemd.com/ophthalmologist.htm>

www.nora.cc

covid.org



Terminology

The 3 subsystems of version, vergence and accommodation must be precisely synchronized for efficient oculomotor control during activities such as reading.

Accommodation	The ability of the eye to change focus from distance to near objects, and is achieved by the ciliary muscle adjusting the shape of the intraocular lens. It occurs reflexively with convergence. Accommodative dysfunction may result in Accommodative Insufficiency, Spasm (overstimulation), or Infacility (slowed)	
Vergence Eye Movements	The simultaneous movement of both eyes, in opposite directions, as needed for single, binocular vision, and stereopsis. 1 st degree fusion is the appreciation of a double image, 2 nd degree fusion is the appreciation of some overlap between the two images, and 3 rd degree fusion is single vision and stereopsis. Stereopsis is binocular depth perception, as the two separate images from each eye are successfully combined /fused into one image in the brain. Vergence eye movements include <i>convergence</i> and <i>divergence</i> .	
Binocular Vision Disorders	Strabismic - eyes are misaligned, may be constant or intermittent <ul style="list-style-type: none"> • Esotropia – eyes turn in • Exotropia – eyes turn out • Hyper tropia – one eye turns up • Hypotropia - one eye turns down 	Non-strabismic <ul style="list-style-type: none"> • Esophoria – eyes have a <i>tendency</i> to turn in • Exophoria – eyes have a <i>tendency</i> to turn out • Hyperphoria – one eye has a <i>tendency</i> to turn up • Hypophoria – one eye has a <i>tendency</i> to turn down
Diplopia	Double vision that suggests misalignment of the eyes. Occurs when the object at which the individual is looking stimulates the fovea (part of the retina that contains the area of most acute vision) of one eye, and a non-foveal part of the retina of the other eye.	
Suppression	A condition in which the visual system ignores the input from one eye, usually associated with strabismus or amblyopia.	
Versional Eye Movements	Saccades - fast, simultaneous eye movements in the same direction, to change the fixation point. During visual scanning, as in reading, the eyes make saccadic movements and stop several times, moving very quickly between each stop. Saccadic dysfunction may include impaired velocity, accuracy, and or initiation of eye movements (ocular motor apraxia). While undershooting is not unusual, overshooting is.	Pursuits - eye movements that allow smooth, continuous viewing of a moving object, and play a significant role in driving and sports. Pursuit dysfunction can include ataxia (jerky quality), and decreased velocity.
Peripheral/Ambient Vision	Allows awareness of position in space, and provides general information needed for balance, movement, coordination, and posture. Unstable ambient vision after BI may cause vertigo, sensitivity to light and motion, and nausea. Also, having intact visual fields does not necessarily correlate with intact peripheral awareness or intact central/peripheral vision integration.	
Vestibulo-Ocular reflex (VOR)	The VOR is one of the fastest reflexes in the human body. At least slight head movement is present all of the time, so the VOR is always important for gaze stabilization. Reflexive eye movements stabilize images on the retina during head movement by producing a compensatory eye movement in the direction opposite to the head movement. In order to have clear vision, head movements must be compensated for almost immediately. Individuals with VOR impairment may complain of vision “looking like a photograph taken with a shaky hand”, or seeing “trails”, dizziness, and imbalance.	




Background

Subtle changes in brain function can occur even after mild TBI/Concussion. Approximately 80% of our perception, learning, cognition and activities are mediated through vision,²³ and greater than half of the circuits in the brain are involved in vision^{3, 53}, including many regions susceptible to shearing during head impacts. Visual system tests probe higher cortical functioning, and assist in detecting functional changes in patients. Abnormal oculomotor and binocular vision skills (ie convergence, accommodation, ocular muscle balance, saccades, and pursuits), vestibular-ocular reflex, and subjective visual complaints are common following TBI, including mTBI^{3,18,19,48,49}. Oculomotor dysfunction has been estimated to be as high as 90% in individuals with TBI, followed by accommodative and convergence deficits^{37,47}. Further, early detection of vision abnormalities may predict individuals at risk of more severe head impacts⁴⁹. Also, it is worth noting that approximately 95% of patients with visual symptoms/complaints caused by TBIs have normal fundoscopic (exam of the retina) findings. An individual can have normal acuity (ie 20/20), but still have impaired oculomotor or eye teaming skills, and therefore impaired functional vision. Also, vision disturbances may be overlooked due to a patient's difficulty in verbally describing impairments.³

Poorer oculomotor function is correlated with more post-concussive symptoms and problems with ADL^{21,48}. Likewise, a recent study by³², found that Convergence Insufficiency was common (~42%) in athletes evaluated within 1 month after a sport-related concussion, and that these athletes had worse neurocognitive impairment and higher symptom scores than did those with normal NPC³². In such cases, more effort will constantly be required for the individual to perform these lower-level actions (ie basic oculomotor control to focus/fuse words) which should be automatic. Subsequently, higher level skills such as comprehension, attention, short-term visual memory, and executive function, will be compromised.^{32,}

⁴⁶ The concept of visual skill hierarchy is illustrated by the Mary Warren Triangle ⁵⁰:



Incidence

Common visual impairments in first year post-TBI¹⁷	<ul style="list-style-type: none"> • Blurred vision • Difficulty with reading • Diplopia/eye strain • Dizziness or disequilibrium in visually crowded environments 	<ul style="list-style-type: none"> • Visual field impairments • Photosensitivity • Color blindness
Incidence specific to mTBI^{17,30,48}	<p>High prevalence in abnormalities of:</p> <ul style="list-style-type: none"> • Saccades • Pursuits • Vergence (most likely convergence) 	<ul style="list-style-type: none"> • Accommodation • Vestibulo-ocular reflex (VOR) • Photosensitivity



Recommendations

(Radomski, 2014). A consensus panel of experts using a nominal group technique has recommended the screening items below. These items have also been confirmed by Fox, et al., 2019⁵³, and have been proposed as “better practice” for vision screening post-TBI with adults, until a validated option becomes available for non-vision specialist clinicians. Items are to be performed in this order.³⁵ *Assessments in italics can also be considered.*

Direct Observation	<ul style="list-style-type: none"> • Direct observation of occupational performance, with attention to behaviors suggesting vision dysfunction. • Also, consider possible time-dependency of symptoms, as they may not manifest immediately. The individual may report complaints only after performing a visual task for certain duration of time.
Questionnaires/ Surveys	<p>Often, patients have difficulty describing their symptoms, and may not recognize them as related to visual deficits (i.e. fatigue, poor attention, dizziness, headache). Using one or more of the following self-report surveys can help the clinician obtain informative answers.³</p> <ul style="list-style-type: none"> • BIVSS (Brain Injury Vision Symptom Survey)^{53, 54} (available at http://links.lww.com/OPX/A248) • COVD QOL outcomes assessment plus photosensitivity interview question ^{12,35} • <i>CISS – score of >21 can indicate impairment</i> ³⁸ • <i>Goodrich, et al. history questions</i>¹⁹ • <i>The Visual Interview</i> ³ <p>Questionnaires if Vestibular Impairment is Suspected</p> <ul style="list-style-type: none"> • <i>The Vestibular Disorders Activities of Daily Living Scale</i>¹⁰ • <i>The Vestibular Activities and Participation Measure</i>^{2,29}
Far Acuity	<ul style="list-style-type: none"> • Chronister Portable Acuity Test ^{35,43} (www.guldenophthalmics.com) • <i>Snellen Acuity Chart</i> <p>Referral to OD is necessary if vision is 20/40 or worse. The numerator refers to the testing distance at which the person recognizes the letter, and the denominator refers to the distance at which the letter being viewed could be identified by a person with normal (20/20) visual acuity.⁴³</p>
Near Acuity/Reading	<ul style="list-style-type: none"> • Chronister Portable Acuity Test reading card^{35,43} <p>Referral to OD is necessary if vision is 20/40 or worse</p>
Accommodation*	<ul style="list-style-type: none"> • Accommodative amplitude testing ⁴³ <ol style="list-style-type: none"> 1. Patch L eye 2. Hold a target (ie, fixation stick) with a 20/30 letter about 1 inch in front of the R eye (it will be blurry at this distance) 3. Slowly move the target away, and ask the patient to report as soon as he or she can identify the target letter 4. Using a ruler, measure the distance from the eye to the target at which the patient was able to identify the letter. Record this measurement in inches 5. Divide 40 by the measurement to determine the amplitude of accommodation. The amplitude should be within 2Diopters of the expected finding (see Addendum A for expected amplitude by age). 6. Patch R eye and repeat.



Recommendations (continued)


Motor Fusion Testing/Eye Alignment Testing*	<ul style="list-style-type: none"> • Near Point of Convergence ^{35,40} <ol style="list-style-type: none"> 1. Use a 20/30 letter target, or pencil tip, placed just above the nose at the brow between the eyes 2. Move the target toward the patient at a rate of about 1-2cm/s, encouraging pt to keep the target single 3. Measure the patient's reported subjective break (target becomes double) in centimeters 4. Then slowly move the target away from the eyes until reported subjective recovery (single again). Measure in centimeters. 5. If the pt does not report diplopia (doubling of the target), the point at which the eyes are observed to lose alignment, and then regain alignment, are recorded as the break and recovery 6. If the eyes are observed to lose alignment and the patient does not report diplopia, this may indicate suppression <p>*Norms – 5cm break and 7 cm recovery</p> <ul style="list-style-type: none"> • Eye Alignment test (options include the cover tests)^{35,43} • <i>Modified Thorington test, using the Bernell Muscle Imbalance Measure card tests (near and far).</i>^{7,51} http://www.bernell.com
Saccades	<ul style="list-style-type: none"> • Developmental Eye Movement test ^{16,35,39} • <i>Northeastern State University College Optometry Oculomotor Test (NSUCO) of saccades</i> ^{27,35} <ul style="list-style-type: none"> *test without corrective lenses* Score is based on patient's ability, accuracy, and the presence of head or body movement during saccadic eye movements. See reference²⁷ for procedure and norms • <i>King-Devick</i> ^{14,49}
Pursuits	<ul style="list-style-type: none"> • NSUCO test of pursuits ^{27,35} <ul style="list-style-type: none"> • Test without corrective lenses. • Score is based on patient's ability, accuracy, and the presence of head or body movement during pursuit eye movements. See reference ²⁷ for procedure and norms
Visual Scanning/Tracking	<ul style="list-style-type: none"> • <i>OSU Visual Tracing test?</i> • <i>Groffman Visual Tracing Test - Designed by Sidney Groffman, OD.,</i> www.bernell.com (no studies)
Visual Fields	<ul style="list-style-type: none"> • Confrontation field testing – finger counting^{1,35} <ul style="list-style-type: none"> • Test without corrective lenses. • This test simultaneously screens for the extinction phenomenon that can accompany unilateral brain damage. See Addendum B.

* Should be trained by a vision specialist before performing fusion tests or accommodative tests.




Visual field impairment and Cranial nerve injury is more likely to occur following moderate or severe TBI.⁴⁸

- mayfieldclinic.com/PE-VisualFieldTest.HTM



Three Cranial Nerves that Supply Innervation to the Six Extraocular Muscles of Each Eye

- ebmedicine.net/topics.php?paaction=showTopicSeg&topic_id=23&seg_id=417

Nerve Palsy	Symptoms	Typical Examination Findings	Differential Diagnosis
Normal	n/a		n/a
Oculomotor (CN III) Palsy	Diplopia, horizontal and vertical		<ul style="list-style-type: none"> • Posterior circulation aneurysm • Brainstem lesion • Microvascular ischemia (if pupil spared) • Cavernous sinus disease
Trochlear (CN IV) Palsy	Torsional (“tilted”) diplopia worse on downward gaze		<ul style="list-style-type: none"> • Brainstem lesion • Posterior circulation aneurysm • Cavernous sinus disease
Abducens (CN VI) Palsy	Horizontal diplopia on lateral gaze to the ipsilateral side		<ul style="list-style-type: none"> • Brainstem lesion • Elevated intracranial pressure • Cavernous sinus disease



Resources for Neuro-Optometric/Sensorimotor Evaluation

OSU College of Optometry

338 W. 10th Ave
Columbus, OH 43210
(614) 292-1113

Galloway Eye Care

Jennifer Mattson, OD
5688 W. Broad St.
Galloway, OH 43119
(614) 853-2020

Riverview Eye Associates

Steven J. Curtis OD, FCOVD
3600 Olentangy River Rd, Unit B
Columbus, OH 43214
(614) 451-7244

Heritage Family Eye Care (primarily pediatric through college age)

Sara Huffman OD, MS
5123 Norwich St., Ste 120
Hilliard, OH 43026
(614) 850-6151

Southwest Eye care

Michelle F. Miller, OD
4140 Hoover Rd.
Grove City, OH 43123
(614) 801-2020

Eyesite of Dublin

Jennifer Mattson, OD
Kimberly Rock, OD
6535 Perimeter Dr.
Dublin, OH 43016
PH (614) 764-7483
FAX (614) 764-7485



Appendix A:
Expected Amplitude of Accommodation by Age⁴³


Age	Expected Number (D)
15	13
18	12
21	11
24	10
27	9
30	8
33	7
36	6
39	5
42	4
45	3
48	2
51	1
54	0

Appendix B:
Rapid Confrontation Screening for Peripheral Visual Field Defects and Extinction

Clinical and Experimental Optometry

Volume 92, Issue 1, pages 45-48, 29 DEC 2008 DOI: 10.1111/j.1444-0938.2008.00280.x
<http://onlinelibrary.wiley.com/doi/10.1111/j.1444-0938.2008.00280.x/full#f1>

Rapid finger-counting confrontation screening requires four responses from the patient (panels A through D). The schematics are shown from the perspective of the clinician, with the patient's eye and covering hand shown in the grey oval. The combinations of fingers shown are examples only, although the clinician must always present either one or two fingers on each hand. A correct response from the patient is always 'two', 'three' or 'four' and never 'one'.



Author: Julie M. Omiatek, MS, OTR/L, CHT

Reviewers: Dr. Steven J Curtis, OD, FCOVD; Karen Rasavage, OTR/L; Michelle F. Miller, OD; Tierney Bumgardner, MOT, OTR/L

Completion date: July 2017

Updated: September 2019

1. Anderson, A. et al. (2009). Rapid confrontation screening for peripheral visual field defects and extinction. Clinical and Experimental Optometry. 92, 45-48.
2. Alghwiri, A. et al. (2012). The development and validation of the Vestibular Activities and Participation Measure. Archives of Physical Medicine and Rehabilitation. 93, 1822-31.
3. Berthold-Lindstedt, et al. (2017). Visual dysfunction is underestimated in patients with acquired brain injury. Journal of Rehabilitation Medicine. 49, 327-332.
4. Brayton-Chung, A. et al. (2016). Action: The role of occupational therapy in concussion rehabilitation. OT Practice. 11-21-2016, 9-12.
5. Capo-Aponte, J. et al. (2012). Visual dysfunctions and symptoms during the subacute stage of blast-induced mild traumatic brain injury. Mil Med. 177(7), 804-813.
6. Cate, Y. & Richards, L., (2000). Relationship between performance on tests of basic visual functions and visual-perceptual processing in persons after brain injury. The American Journal of Occupational Therapy. 54(3), 326-334.
7. Cebrian, J. et al. (2014). Repeatability of the modified Thorington card used to measure far heterophoria. Optom Vis Sci. 91(7). 786-92.
8. Ciuffreda, KJ. (2002). The scientific basis for and efficacy of optometric vision therapy in nonstrabismic accommodative and vergence disorders. Optometry. 73, 735-762.
9. Ciuffreda, KJ, et al. (2008). Vision therapy for oculomotor dysfunctions in acquired brain injury: A retrospective analysis. Optometry. 79, 18-22.
10. Cohen, H. (2014). Use of the Vestibular Disorders Activities of Daily Living Scale to describe functional limitations in patients with vestibular disorders. Journal of Vestibular Research. 24, 33-38.
11. Cooke, J and Zacko, C. (2014) Post-concussion Syndrome, Persistent Symptomatic Concussion, Related Sequelae, and Treatment of Mild Closed Head Injury (ch 22). Concussions in Athletics: From Brain to Behavior. Springer Science + Business Media, New York
12. Daugherty, K. et al. (2007). Evaluating changes in quality of life after vision therapy using the COVD Quality of Life Outcomes Assessment. Optometry and Vision Development. 38, 75-81.
13. Finn, C. & Waskiewicz, M. (2015). The role of occupational therapy in managing Post-Concussive Syndrome, published by the American Occupational Therapy Association, Inc., Physical Disabilities Special Interest Section Quarterly, 38(1), 1-4.
14. Galetta, K.M., et al. (2011). The King-Devick test and sports-related concussion: study of a rapid visual screening tool in a collegiate cohort. Journal of the Neurological Sciences. 309, 34-39.
15. Gallaway, M. et al. (2017). Vision Therapy for Post-Concussion Vision Disorders. Optom Vis Sci. 94(1), 68-73.
16. Garzia, R. et al. (1990). A new visual-verbal saccade test: The Developmental Eye Movement test (DEM). Journal of the American Optometric Association. 61(2), 124-135.
17. Greenwald, B. D., Kapoor, N., & Singh, A. D. (2012). Visual impairments in the first year after traumatic brain injury. Brain Injury, 26, 1338-1359.
18. Goodrich, G., et al. (2007). Visual function in patients of a polytrauma rehabilitation center: A descriptive study. Journal of Rehabilitation Research & Development. 44(7), 929-936.
19. Goodrich, G. et al. (2013). Development of a mild traumatic brain injury-specific vision screening protocol: A Delphi study. JRRD. 50(6). 757-768.
20. Han, Y., et al. (2004). Reading-related oculomotor testing and training protocols for acquired brain injury in humans. Brain Research Protocols. 14, 1-12
21. Heitger, M. et al. (2009). Impaired eye movements in post-concussion syndrome indicate suboptimal brain function beyond the influence of depression, malingering, or intellectual ability. Brain. 132; 2850-2870.
22. Hoyt, CS., (2007). Brain injury and the eye. Eye. 21, 1285-1289.
23. Kelts, E. (2010). The basic anatomy of the optic nerve and visual system (or, why Thoreau was wrong). Neurorehabilitation. 27, 217-222.
24. Kelts, E. (2010). Traumatic brain injury and visual dysfunction: A limited overview. Neurorehabilitation. 27, 223-229.



25. Kapoor, N., et al. (2004). Oculomotor rehabilitation in acquired brain injury: A case series. Archives of Physical Medicine and Rehabilitation. 85, 1667-1678.
26. Lavrich, JB. (2010). Convergence insufficiency and its current treatment. Current Opinion in Ophthalmology. 21(5), 356-360.
27. Maples WC. NSUCO Oculomotor test. Santa Ana, CA: Optometric Extension Program. 1995.
28. Marshall, S. et al. (2015). Updated clinical practice guidelines for concussion/mild traumatic brain injury and persistent symptoms. Brain Injury. 29:6, 688-700.
29. Mueller, M. et al. (2015). Subscales of the Vestibular Activities and Participation questionnaire could be applied across cultures. Journal of Clinical Epidemiology. 68, 211-219.
30. Master, CL et al. (2015). Vision diagnoses are common after concussion in adolescents. Clinical Pediatrics. July 7.
31. Padula, W. & Argyris, S. (2015). Post Trauma Vision Syndrome: Part I and Part 2, Traumatic Brain Injury Resource Guide for the Centre for Neuro Skills.
32. Pearce, K. et al. (2015). Near Point of Convergence after a sport-related concussion: Measurement reliability and relationship to neurocognitive impairment and symptoms. American Journal of Sports Medicine. 43(12), 3055-3061.
33. Politzer, T. & Lenahan, D. (2010). Double vision caused by neurologic disease and injury. Neurorehabilitation. 27, 247-254.
34. Padula, W. et al. (2008). The need for optometric rehabilitation for our veterans who have incurred traumatic brain injury – Senate Bill 1999/House Bill 3558. Optometry-Journal of the American Optometric Association - Guest Editorial. 170-171.
35. Radomski, M. et al. (2014) Composition of a vision screen for service members with Traumatic Brain Injury: Consensus using a modified nominal technique. American Journal of Occupational Therapy. July/August 2014, Vol. 68, 422-429.
36. Ripley, D. et al. (2010). The Vision Clinic: An interdisciplinary method for assessment and treatment of visual problems after Traumatic Brain Injury. Neurorehabilitation. 27, 231-235.
37. Roberts, P. et al. (2016). A conceptual model for vision rehabilitation. Journal of Rehabilitation Research & Development (JRRD). 53(6). 693-704.
38. Rouse, M et al. (The CITT Investigator Group). (2009) Validity of the CISS: A confirmatory study. Optom Vis Sci. 86(4). 357-63.
39. Richman, J.E. & Garzia, R. P. (2015). The Developmental Eye Movement test, version 2.5. South Bend, IN: Bernell
40. Scheiman, M. et al. (2003). Nearpoint of Convergence: Test procedure, target selection, and normative data. Optometry and Vision Science. 80(3). 214-225.
41. Scheiman, M & Wick, B (2007). Clinical Management of Binocular Vision: Heterophoric, Accommodative, and Eye Movement Disorders, 4th ed., Wolters Kluwer.
42. Scheiman, M. et al. (2010). Vision therapy/orthoptics for symptomatic convergence insufficiency in children: Treatment kinetics. Optom Vis Sci. 87(8), 593-603
43. Scheiman, M. et al. (2011). Understanding and Managing vision deficits: A guide for occupational therapists (3rd ed). Thorofare, NJ: Slack.
44. Shin, H. et al. (2011). Effectiveness of vision therapy for convergence dysfunctions and long-term stability after vision therapy. Ophthalmic & Physiological Optics. 31, 180-189.
45. Thiagarajan, P. et al. (2011). Vergence dysfunction in mild traumatic brain injury (mTBI): a review. Ophthalmic Physiol Opt. 31: 456-68.
46. Thiagarajan, P. et al. (2014). Oculomotor neurorehabilitation for reading in mild traumatic brain injury (mTBI): An integrative approach. NeuroRehabilitation. 34:129-146.
47. Tjarks, B. et al. (2013). Comparison and utility of King-Devick and ImPACT composite scores in adolescent concussion patients. Journal of the Neurological Sciences. (Article in Press)
48. Ventura, R. et al. (2014). The neuro-ophthalmology of head trauma. Lancet Neurology. 13: 1006-16.
49. Ventura, R. et al. (2015). The Concussion Toolbox: The role of vision in the assessment of concussion. Seminars in Neurology. 35, 599-606.
50. Warren, M. (1993). A hierarchical model for evaluation and treatment of visual perceptual dysfunction in adult acquired brain injury, Part 1. The American Journal of Occupational Therapy. 47(1), 42-54.
51. Wong, E. et al. (2002). Interexaminer repeatability of a new, modified prentice card compared with established phoria tests. Optom Vis Sci. 79(6). 370-5.
52. Zoltan, B. (2007). Vision, perception, and cognition: A manual for the evaluation and treatment of the adult with acquired brain injury, 4th ed. Thorofare, NJ: SLACK, Inc.



53. Fox, S. et al. (2019). Vision Rehabilitation after Traumatic Brain Injury. Physical Medicine and Rehabilitation Clinics of North America. 30, 171-188.
54. Laukkanen, H. et al. (2017). Brain Injury Vision Symptom Survey (BIVSS) Questionnaire. Optometry and Vision Science. 94(1), 43-50.
55. Armstrong, RA. (2018). Visual Problems Associated with Traumatic Brain Injury. Clinical and Experimental Optometry. 101, 716-726.

